

Malaria vaccines indicator

Values and preferences

Perception of the target population on the risk of disease

General awareness of malaria risk is high in endemic settings, with many communities recognizing malaria as common, severe, and life-threatening, especially for young children and pregnant women. ¹ Peer-reviewed articles demonstrate the perception of malaria among caregivers of young children in malaria endemic countries in sub-Saharan Africa. ^{2, 3, 4}

Sources

1.

Abdul Rahim FA, Mahmud MAF, Abdul Mutalip MH, Yoep N, Aminuddin MAH et al. A scoping review of community knowledge in malaria prevention and control programmes. PLoS One. 2025;20(7):e0328703. doi:10.1371/journal.pone.0328703.

2.

Ojakaa DI, Ofware P, Machira YW, Yamo E, Collymore Y et al. Community perceptions of malaria and vaccines in the South Coast and Busia regions of Kenya. Malar J. 2011;10:147. doi:10.1186/1475-2875-10-147.

3.

Bingham A, Gaspar F, Lancaster K, Conjera J, Collymore Y et al. Community perceptions of malaria and vaccines in two districts of Mozambique. *Malar J.* 2012;11:394. doi:10.1186/1475-2875-11-394.

4.

Yaya Bocoum FIK, Kouanda S, Hinson L, Collymore Y, Ba-Nguz A et al. Community perceptions of malaria vaccines: qualitative research from the sanitary districts of Kaya and Hounde in Burkina Faso. *Glob Health Promot.* 2014;21(1):76–87. doi:10.1177/1757975913507729.