

Table 2. Vaccines used in the Expanded Program on Immunization (EPI) over time

1977-1981	1982-1986	1987-1991	1992-1996	1997-2001	2002-2007	2008-2009
BCG	BCG	BCG x 2	BCG	BCG	BCG	BCG
DTP x 2	DTP x 3	DTP x 4	DTP x 5	DTP x 5	DTP x 5	DTP x 2
OPV x 2	OPV x 3	OPV x 4	OPV x 5	OPV x 5	OPV x 5	OPV x 5
	Measles	Measles	Measles	Measles	Measles	Measles
	Rubella	Rubella	Rubella x 2	MMR	MMR	MMR
			HB x 3	HB x 3	HB x 3	DTP-HB x 3
			JE x 2	JE x 2	JE x 3	JE x 3
Typhoid*	Typhoid*	Typhoid*	-	-	-	Influenza
	TT x 2	TT x 2	TT x 3	TT x 3	dT x 3	dT x 3

*killed whole-cell parental vaccine

BCG = bacille Calmette-Cuérin; OPV=Oral Poliovirus Vaccine; HB=Hepatitis B vaccine; DTP=Diphtheria-Tetanus-Pertussis vaccine; MMR=Mump-Measles-Rubella vaccine; JE=Japanese Encephalitis vaccine; dT=Diphtheria-Tetanus vaccine for adolescent and adults. The number following “x”= the total doses of vaccines per series.